

**Effects of Clomiphene citrate on the histological structure  
of the Iraqi domestic chicken's ovaries**

**Hayfaa A Al-Shammary**

**Department of Medical Analysis, College of Science,  
University of Thi Qar, Thi Qar, Iraq**

**Abstract**

Clomiphene citrate is commonly used to induce ovulation in women. However, the effects of clomiphene citrate on the domestic chickens ovary's have not been fully elucidated. Therefore, we used histological analyses to examine the ovaries of Iraqi domestic chickens administered clomiphene citrate (25 mg/kg of body weight) daily for 30 and 60 days. Our results demonstrated that the ovaries from the hens that were administered clomiphene citrate for 30 and 60 days were not morphologically or histologically distinct. However, compared to the control group, both these groups showed a significant treatment effect on ovarian diameter and the number of follicles ( $P < 0.05$ ), from the result we can conclusion that the clomophene citrate stimulating the ovalation in domestic chickens .

**Keywords:** Clomiphene citrate, chicken ovary, histological structure.

**Introduction**

Clomiphene citrate, commonly used to induce ovulation in women, is an anti-estrogenic agent that stimulates the secretion of pituitary gonadotrophic hormones. It stimulates ovulation by blocking the effect of estrogens at the receptor sites in the hypothalamus and pituitary gland (Kerin etal ,1985). The structure of clomiphene is similar to that of estrogen. Similar to other hormones, estrogen exerts its effects throughout the body by occupying its receptors; the occupied receptor signals the organ. In the brain, estrogen directs the release of follicle-stimulating hormone (FSH) (reference). Clinical studies have demonstrated that while clomiphene citrate induces ovulation in 80% of physically capable

women, it results in a pregnancy rate of only 40% (Speroff *et al.*, 1983). O'Keefe and Marrone (1986) reported that clomiphene citrate inhibits androgen and estrogen production in thecal cells in vitro by decreasing the activity of C17–C20 lyase and aromatase. The effects of clomiphene citrate on the hypothalamic-pituitary axis are widely accepted.

Clomiphene citrate antagonizes estrogen receptor (ESR) signaling in female reproductive tissues including the ovaries in rodents (Chaube *et al.*, 2006; London *et al.*, 2000), fallopian tubes in chickens (Sutherland *et al.*, 1977), and endometrium in humans (Gonen and Casper 1990; Homburg *et al.*, 2006). It also acts as a strong agonist in non-reproductive tissues such as bone and the liver (Turner *et al.*, 1998). In rats, clomiphene citrate induces apoptosis of ovarian granulosa cells, which can be rescued by 17 $\beta$ -estradiol (E2) in vivo (Chaube *et al.*, 2006, Nagao and Yoshimura, 2001). Thus, clomiphene citrate acts as an ESR antagonist that regulates apoptosis in the tissue and cellular targets of estrogen signaling. Apoptotic changes in human chorionic villi and decidual tissues may increase the risk of tubal ectopic pregnancy (Kokawa *et al.*, 1998).

Clomiphene citrate may impair fertility by affecting the cervical mucus and causing endometrial dysfunction. The drug is as efficacious at a dose of 50 mg as it is at a dose of 100 mg; moreover, the lower dose helps avoid some undesirable side effects. However, other adverse effects may occur at this dose, including multiple pregnancies, an increase in the rate of multiple births, and ovarian hyperstimulation. Ovarian cancer may also develop owing to 50 mg clomiphene citrate administration; however, this claim is yet to be substantiated (Sovino *et al.*, 2002).

The aim of the present study was to assess the effects of chronically administered clomiphene citrate on ovulation in domestic chickens. We used the left ovary and oviduct to represent the reproductive organs of the hen. The ovary consists of an outer cortex enveloping a vascular medulla and containing ovarian follicles of various sizes. Covering the cortex is a layer (germinal epithelium) of cuboidal or flattened cells. Below the epithelium is the tunica albuginea, composed of

dense connective tissue, and below the tunica albuginea is a stroma of loose connective tissue (Bacha and Bacha, 2000).

## **Materials and Methods**

Thirteen Iraqi domestic hens, *Gallus gallus domesticus*, aged 30 weeks were subjected to a daily 14-h light to 10-h dark cycle. The hens were provided water and commercially available chicken feed. They were treated daily for 30 and 60 days with clomiphene citrate (25 mg/kg of body weight). After killing the hens, the ovaries were removed, fixed in 10% formalin, dehydrated, and embedded in paraffin. Sections (5 mm) were prepared and stained with hematoxylin and eosin (H & E; Luna, 1960). The histological structure of the ovary was examined using light microscopy, and the data were recorded and analyzed. Data are expressed as mean  $\pm$  standard error of the mean (SEM) and analyzed by two-way analysis of variance followed by Fisher's least significant difference analysis. Differences among means were considered significant at  $P < 0.05$ . Analyses were performed using Statistical Product and Service Solutions, 2006, software (IBM Corporation, Armonk, NY, USA).

## **Results**

No significant histopathological differences were observed in the ovaries of hens following administration of clomiphene citrate (25 mg/kg of body weight) daily for 30 or 60 days in comparison with those of the control group (Fig. 1, 2, and 3). However, histometric evaluation showed that clomiphene citrate administration for 30 and 60 days stimulated ovulation and significantly affected the diameter of the follicles, compared to the control group ( $P < 0.05$ ). The diameter of the ovarian follicles in the group of hens treated with clomiphene citrate during all stages of follicular maturation (follicle selection, slow-growing white follicles, and final differentiation) was similar at 30 and 60 days of treatment but significantly greater than that for hens in the control group, as shown in Table 1. While there was no significant difference in the number of follicles (10 follicles in 15 mm<sup>2</sup>) between 30 and 60 days of treatment, administration of clomiphene citrate significantly increased the

number of ovarian follicles in both these groups compared to the number in the control hens (6 follicles in 15 mm<sup>2</sup>, P < 0.05) (Fig. 4, 5, and 6).

We observed numerous small immature follicles on the surface of the stalks in the cortical tissue (Fig. 7) as well as on the surface of mature large follicles (Fig. 8 and 9). There was no histopathological effect of clomiphene citrate on the layers of the ovaries between the two experimental groups (30 and 60 days of treatment), which is consistent with an increase in the yolk-laden oocytes with rounded nuclei (germinal vesicles). The oocytes were surrounded by several layers. Large follicles were suspended from the surface of the ovary by stalks of cortical tissue comprising follicle of varying sizes (Fig. 7). The surface of many large follicles had one or two small immature follicles (Fig. 8).

**Table 1. Diameter of ovarian follicles following daily chronic administration of clomiphene citrate to domestic hens.**

Period	30 days	60 days	
	Diameter (mm <sup>2</sup> )	Diameter (mm <sup>2</sup> )	
<b>Control</b>	5.0 ± 0.543 a	5.10 ± 0.543a	5.05±0.53
<b>Clomiphene citrate(25mg/kgbody weight)</b>	10.7 ± 0.232 b	10.1 ± 0.248 b	10.4±0.23
<b>Mean</b>	7.85±0.44	8.05±0.540	
<b>Interaction LSD</b> 4.02			

\*Values represent mean ± SEM. Values with nonidentical superscripted letters (a–b) are considered significantly different P < 0.05.




Fig. 1. Tissue from a hen in the control group showing ovarian follicles (arrows). H & E stain, magnification  $\times 400$ .




Fig. 2. Tissue from a hen administered clomiphene citrate (25 mg/kg body weight) daily for 30 days showing the various sizes of ovarian follicles (arrows). H & E stain, magnification  $\times 400$ .






Fig. 3. Tissue from a hen administered clomiphene citrate (25 mg/kg body weight) daily for 60 days showing the various sizes of ovarian follicles (arrows). H & E stain, magnification  $\times 400$ .




Fig. 4. Tissue from a hen in the control group showing numerous ovarian follicles (arrows). H & E stain, magnification  $\times 170$ .




Fig. 5. Tissue from a hen administered clomiphene citrate (25 mg/kg body weight) daily for 30 days showing numerous ovarian follicles (arrows). H & E stain, magnification  $\times 170$ .




Fig. 6. Tissue from a hen administered daily clomiphene citrate (25 mg/kg body weight) for 60 days showing numerous ovarian follicles (arrows). H & E stain, Magnification  $\times 170$ .




Fig. 7. Image taken during morphological analysis showing the growth of numerous ovarian follicles on the surface of ovarian stalks (arrows). H & E stain,30 x.



Fig. 8. Image taken during morphological analysis showing small follicles on the surface of mature follicles (arrows).






Fig. 9. Image taken during morphological analysis showing numerous immature follicles on the surface of one follicle (arrows), 30 x .

## **Discussion**

Our results demonstrated that the ovaries from Iraqi domestic chickens administered clomiphene citrate (25 mg/kg of body weight) daily for 30 days and 60 days were not morphologically or histological distinct. However, we found a significant effect of treatment for both of these groups in comparison with hens in the control group on ovarian diameter and on the number of follicles.

Clomiphene citrate exhibits estrogenic and antiestrogenic effects and induces ovulation in a complex fashion. Therefore, clomiphene citrate could have a direct estrogenic effect on gonadotrophs, enhancing sensitivity to gonadotropin-releasing hormone (GnRH). Moreover, clomiphene has a direct estrogenic effect on the ovary, which is similar to that described for the pituitary gland. It sensitizes the granulosa cells in the follicle to the action of gonadotrophins and upregulates aromatase activity. The agonistic or antagonistic effects of clomiphene citrate on the

effectors in the genital tract may also be due to different populations of  $\alpha$  and  $\beta$  estrogen receptors in those tissues (Goldstein *et al.*, 2000). Adashi (1984) showed that there are three sites of action for clomiphene citrate. The primary site is in the hypothalamus, where clomiphene citrate blocks the negative feedback of estrogen, thereby enhancing the release of gonadotrophins. The secondary sites are in the pituitary, where clomiphene citrate enhances the GnRH-stimulated release of FSH, and in the ovary, where its effects have not been fully elucidated.

Clomiphene citrate inhibits progesterone synthesis in rat follicles (Laufer *et al.*, 1982) and granulosa cells (Welsh *et al.*, 1984), in monkey luteal cells (Westfahl and Resko, 1983), and in hen granulosa cells (Sgarlata *et al.*, 1984). O'Keffe and Marrone (1987) conducted an in vitro study to show that clomiphene citrate inhibits luteinizing hormone-stimulated androstenedione and estrogen production in a dose-dependent manner in hen theca cells.

Shao *et al.* (2009) showed that during chronic clomiphene citrate administration in rats, isthmus-specific apoptosis of epithelial cells and activation of ESR2 $\alpha$  in cilia act in parallel to block gamete and embryo passage through the fallopian tube, eventually resulting in tubal ectopic pregnancy. Kokawa *et al.* (1989) showed that the apoptotic changes in human chorionic villi and decidual tissues may increase the risk of tubal ectopic pregnancy. However, clomiphene citrate causes ovarian and uterine abnormalities (Nagao and Yoshimura, 2001) and may have adverse effects on the fallopian tube or on apoptotic signaling pathways in this tissue (Kokawa *et al.*, 1998).

In this study we are find that the clomiphene citrate had a good effect on hens ovaries which causes significantly increase of number and diameters of the follicles and increased early egg production .

## **References**

Adashi EY. (1984). Clomiphene citrate: mechanism(s) and site(s) of action; a hypothesis revisited. *Fertil. Steril.* 42:331–344.

Asem EK, Hertelendy F. (1985). Antiestrogens inhibit the cholesterol side-chain cleavage enzyme activity. *Biol. Reprod. Suppl.* 1, 32, Abstr. 105.

Asem EK, Hertelendy F. (1986). Clomiphene and tamoxifen inhibit the cholesterol side-chain cleavage enzyme activity in hen granulosa cells. *J. Reprod. Fert.* 77:153–158.

Bacha WJ, Bacha ML. (2000). *Color Atlas of Veterinary Histology*. 2nd Ed. Chaube SK, Prasad PV, Tripathi V, Shrivastav TG. (2006). Clomiphene citrate inhibits gonadotropin-induced ovulation by reducing cyclic adenosine 30,50-cyclic monophosphate and prostaglandin E2 levels in rat ovary. *Fertil. Steril.* 86(Suppl 4):1106–1111.

Goldstein SR, Siddhanti S, Ciaccia AV, Plouffe L Jr. (2000). A pharmacological review of selective oestrogen receptor modulators. *Hum. Reprod.* 6:212–224.

Gonen Y, Casper RF. (1990). Sonographic determination of a possible adverse effect of clomiphene citrate on endometrial growth. *Hum. Reprod.* 5:670–674

Homburg R, Pap H, Brandes M, Huirne J, Hompes P, Lambalk CB. (2006). Endometrial biopsy during induction of ovulation with clomiphene citrate in polycystic ovary syndrome. *Gynecol. Endocrinol.* 22:506–510.

Kerin JF, Liu JH, Phillipou G, Yen SS. Evidence for a hypothalamic site of action of clomiphene citrate in women. *J Clin Endocrinol Metab* 1985;61:265–8.

Kokawa K, Shikone T, Nakano R. (1998). Apoptosis in human chorionic villi and decidua in normal and ectopic pregnancy. *Mol. Hum. Reprod.* 4:87–91.

Laufer N, Reich R, Braw R, Shenker JG, Tsafiriri A. (1982). Effect of clomiphene citrate on preovulatory rat follicles in culture. *Biol. Reprod.* 11:463–471.

London SN, Young D, Caldito G, Mailhes JB. (2000). Clomiphene citrate-induced perturbations during meiotic maturation and cytogenetic abnormalities in mouse oocytes in vivo and in vitro. *Fertil. Steril.* 73:620–626.

Luna LG. (1960). Manual of Histological Staining Methods of the Armed Forces Institute of Pathology. 3rd ed. by McGraw Book Company. New York, London.

Macut D, Micic D, Suvajdzic N, Sumarac M, Kendereski A, Zoric S, Elezovic I, Petrovic R, Cvijovic G, Gligorovic P. (2000). Ovulation induction and early pregnancy loss in a woman susceptible to autoimmune diseases: a possible interrelationship. *Gynecol. Endocrinol.* 14:153–157.

Nagao T, Yoshimura S. (2001). Oral administration of clomiphene to neonatal rats causes reproductive tract abnormalities. *Teratog. Carcinog. Mutagen.* 21:213–221.

O'Keefe, RK, Marrone BL. (1986). Androgen and oestrogen production by clomiphene citrate in avian theca cells. *J. Reprod. Fertil.* 78:541–548.

Onsperoff L, Glass RH, Kase NG. (1983). *Clinical Gynecologic Endocrinology and Infertility*, pp. 523–544. Williams and Wilkins, Baltimore.

Schmierer B, Schuster MK, Shkumatava A, Kuchler K. (2003). Activin and follicle-stimulating hormone signaling are required for long-term culture of functionally differentiated primary granulosa cells from the chicken ovary. *Biol. Reprod.* 68:620–7.

Sgarlata, CS, Mikhail G, Hertelendy, F. (1984). Clomiphene and tamoxifen inhibit progesterone synthesis in granulosa cells: comparisons with estradiol. *Endocrinology.* 114:2032–2038.

Shao R, Egecioglu E, Weijdegard B, Kopchick JJ, Fernandez-Rodriguez J, Andersson N, Billig H. (2007). Dynamic regulation of estrogen receptor-alpha isoform expression in the mouse fallopian tube: mechanistic insight into estrogen-dependent production and secretion of insulin-like growth factors. *Am. J. Physiol. Endocrinol. Metab.* 293:E1430–E1442.

Sovino H, Sir-Petermann T, Devoto L. (2002). Clomiphene citrate and ovulation induction. *Reproductive BioMedicine.* 4(3):303–310.

Speroff L, Glass RH, Kase NG. (1983). *Clinical Gynecologic Endocrinology and Infertility*, pp. 523-544. Williams and Wilkins, Baltimore

Sutherland R, Mester J, Baulieu EE. (1977). Tamoxifen is a potent “pure” antioestrogen in chick oviduct. *Nature*. 267:434–435.

Turner RT, Evans GL, Sluka JP, Adrian MD, Bryant HU, Turner CH, Sato M. (1998). Differential responses of estrogen target tissues in rats including bone to clomiphene, enclomiphene, and zuclomiphene. *Endocrinology*. 139:3712–3720.

Welsh TH, Jia X, Jones PBC, Zhuang L, Hsueh AJW. (1984). Disparate effects of triphenylethyleneantiestrogens on estrogen and progesterin biosynthesis by cultured rat granulosa cells. *Endocrinology*. 115:1275–1282.

Westfahl PK, Resko JA. (1983). Effects of clomiphene on luteal function in the nonpregnant cynomolgus macaque. *Biol. Reprod*. 29:963–969.

## تأثير عقار كلوميفين سترات على التركيب النسيج لمبايض الدجاج المحلي العراقي

هيفاء عبد علي الشمري

قسم التحليلات المرضية كلية العلوم جامعة ذي قار

يستخدم عقار كلوميفين سترات عادة للحث على الإباضة عند النساء. ان تأثير عقار كلوميفين سترات على مبايض الدجاج لم توضح تماما لذلك اجريت هذه الدراسة النسيجية لفحص تأثير عقار كلوميفين سترات بجرعتين (25 ملغم / كغم من وزن الجسم) يوميا لمدة 30 و 60 يوما على التركيب النسيجي للمبايض. أظهرت النتائج عدم وجود اي تأثيرات مرضية نسيجية لمبايض الطيور المعالجة بالجرعتين من عقار كلوميفين سترات لمدة 30 و 60 يوما ، كما وأظهرت النتائج ان هذه الجرعة لها تأثير معنوي عند مستوى احتمال ( $P > 0.05$ ) على قطر وعدد بصيلات المبيضية. من هذه النتائج نستنتج بأن عقار كلوميفين سترات يحفز التبويض في مبايض الدجاج المحلي.