

**Detection of Fibronectin Binding Protein (*fnb A*) gene in
Staphylococcus aureus isolates**

<https://doi.org/10.32792/utq/utj/vol14/1/2>

**Abbas D. Mater¹, Zainab Dakhil Degaim², Abdulazeez
Salih Abdulazeez¹**

¹ College of Veterinary Medicine / University of Thi-Qar,

² College of Medicine / University of Thi-Qar,

Email: Zainab-d@utq.edu.iq

Email: Dr.abbas-mater@utq.edu.iq

Abstract:

This study designed to molecular evaluation of the fibronectin-binding proteins gene (*fnb A*) in clinical isolates of *S. aureus* which recovered from 120 swabs from pharyngitis and tonsillitis patients during the period from February, 2017 to November, 2017. Among all bacterial isolates, only 32 (26.66%) were positive on mannitol salt agar and biochemical methods, which identified as *S. aureus*. The molecular analysis of *fnb A* gene was done by employing the polymerase chain reaction (PCR). The results revealed that 59% of isolates were found positive for *fnb A* gene.

الخلاصة

صممت هذه الدراسة للتقييم الجزيئي لجين (البروتينات المرتبطة بالفايبرونكتين) في عينات بكتريا المكورات العنقودية الذهبية السرييرية المعزولة من ١٢٠ مسحة قطنية من مرضى التهاب البلعوم والتهاب اللوزتين خلال الفترة من شباط، ٢٠١٧ الى تشرين الثاني، ٢٠١٧. اثنان وثلاثون عزلة (٢٦,٦٦%) من كل البكتريا المعزولة شخصت على انها بكتريا المكورات العنقودية الذهبية بالاعتماد على النمو على وسط اكار المنترول الصلب والاختبارات البايوكيميائية، اوضحت نتائج التحليل الجزيئي باستخدام تقنية تفاعل البلمرة الجزيئي ان ٥٩% من عينات المكورات العنقودية الذهبية تحتوي على جين.

Keyword: *S. aureus*, PCR, *fnb A* gene.

Introduction:

Staphylococcus aureus is a successful commensal bacteria that efficiently colonizes both human and hosts (Lowy, 2011). *S. aureus* is an

opportunistic bacteria with its capability to occupy and persist in unethical phagocytes: osteoblasts, fibroblasts and diverse types of epithelial cells. The infectious potential of this bacteria detected by a huge number of cell related and extracellular virulence factors, some of which are concerned in the adhesion process and others in the bacterial invasion (Holban *et al.*, 2013). The intracellular presence of *S. aureus* in host tissues has been recently highlighted, advising the ability of the pathogen to dodge the host defense mechanisms and to persist in host cells (Zautner *et al.*, 2010).

The adherence of *S. aureus* to human tissue was mediated by a number of genes encoding of factors related to cell wall microbial surface components, such as microbial surface components recognizing adhesive matrix molecules (MSCRAMMs), this molecules play an significant role in *S. aureus* pathogenesis, allowing its transition from the bacteremia stage to infections such as endocarditis or osteomyelitis (Bur *et al.*, 2013). The MSCRAMMs group includes protein A, the fibronectin-binding proteins (*fnb* A and B), collagen-binding protein (*cna*), elastin-binding protein, clumping factors A and B (*ClfA* and *ClfB*), those virulence elements are significant for the invasiveness and development of infections by invasive *S. aureus* (Sabat *et al.*, 2006; Sivaraman *et al.*, 2009).

The invasive infections caused by *S. aureus* isolated strains, which harbor the *fnbA* and *fnbB* adhesins genes (Peacock *et al.*, 2000), and those virulence factor used to prevent the phagocytosis process by leucocytes (Higgings *et al.*, 2006). The present study aimed to investigate the prevalence of *S. aureus* in pharyngitis and tonsillitis patients, PCR-based technique was utilized to detect the gene encoding for the fibronectin-binding protein (*fnb* A).

Material and Methods:

Ethical approval

This research was approved by the Veterinary Medicine College Ethics Committee, Thi-Qar University, Thi-Qar Province, Iraq.

Laboratory methods

Totally, *S. aureus* isolates were isolated from 120 swabs which collected from pharyngitis and tonsillitis patients whom admitted to ENT unit in AL-Habbuby Teaching Hospital of Thi-Qar province through the period from February, 2017 to November, 2017 and identified depending on cultural properties on mannitol salt agar and blood agar (LAB/ United Kingdom), followed by biochemical tests (Harley and Prescott, 2002). The confirmed diagnosis was performed by using API system (BioMerieux/France). Staphylo Monotec test kit Plus (Mast/United Kingdom) used to identify all *S. aureus* isolates as serological diagnosis of these bacteria.

Extraction of bacterial DNA

All isolates of *S. aureus* were inoculated on Nutrient broth (NB) (LAB/United Kingdom) and incubated for 24h at 37°C. Bacterial DNA was extracted from fresh overnight culture in NB broth by using DNA Bacteria plus kit (Geneaid/Korea) according to the manufacturer's instructions.

PCR diagnosis of *fnb A* gene

The specific primer pairs of *fnb A* gene as following: forward: 5'- ATC AGC AGA TGT AGC GGA AG -3' and reverse: 5'- TTT AGT ACC GCT CGT TGT CC -3'. The PCR cycling conditions of *fnb A* gene: initial denaturation at 94°C for 2 min, followed by 30 cycles of denaturation at 94°C for 1min, annealing at 55°C for 1min, extension at 72°C for 1min and final extension for 5min after the last cycle (Abraham, and Jefferson, 2010) with modification. Electrophoresis of PCR product was carried out in 1.4% agarose gel and the presence of a 198 bp band indicate a positive result for *fnb A* gene.

Results and Discussion:

From of 120 collected swabs, only 32 (26.66%) was identified as *S. aureus*. The *S. aureus* was noticeable causer of pharyngitis and tonsillitis infection but create less often when compared to the *Streptococcus pyogenes* pharyngitis (Jeong *et al.*, 2007; Gowrishankar *et al.*, 2013). The current results were disagreement with results of local study performed by Dakhil and Hamim, (2016) showed that the occurrence of *S. aureus* was 64.58%. The percentage of *S. aureus* was higher than the

results of study done by Sadoh *et al.*, (2008) indicated that the occurrence of this agent was 12.83%. The present data were differenced from results of studies performed by Al-Ahmary *et al.*, (2012) and Thomas, (2012) demonstrated that a percentage of *S. aureus* was 44% and 52% respectively.

Among the examined *S. aureus*, 19 amplified of the targeted gene, the percentage was 59% with the molecular weight of approximately 198 bp (Fig.1). The present study disagreed with results of Duran *et al.*, (2010) revealed nearly all of the strains were positive for *fnb A* gene (97.7%). While, Nashev *et al.*, (2004) noted that all *S. aureus* isolates harbored the goal gene.

On other hand, the prevalence of *fnb A* gene in recent study was slightly with the results of other studies, like Mirzaee *et al.*, (2015) and Gowrishankar *et al.*, (2016) showed that high percentage of methicillin resistant *S. aureus* had *fnbA* gene, were 77.8 and 82.2%, respectively. While, the present results were incorporated with results of study performed by Kiavari *et al.*, (2012) recorded that *fnb* was identified in 65.29% of clinical and 54.67% nasal *S. aureus* strains and had higher prevalence in isolates from endocarditis, osteomyelitis and burn associated wounds.

Interestingly, some studies recorded that high percentages of *S. aureus* isolates harbored *fnbA* gene whereas no detectable in MRSA isolates like Tangchaisuriya *et al.*, (2014) documented 93.5% of *S. aureus* had the current gene and lacked in all MRSA isolates. Also Taneike *et al* (2006) described that completely isolates of MRSA which remote from nosocomial eruptions in Japan lacked *fnbB*. While, the results of Wisniewska *et al.*, (2008) showed that *fnbA* and *cna* established in 90% and 63%, respectively of all MRSA strains. O'Neill *et al.*, (2007) showed that a higher percentage of methicillin sensitive *S. aureus* (14%) than MRSA (0%) harbored the goal gene.




Fig.(1): Agarose gel electrophoresis of *fnb A* gene amplification, M: ladder, 1,3, 5-7 : positive results, 2,4,Negative result.

Reference:

- ◆ Abraham, N M. and Jefferson, K K. (2010). A low molecular weight component of serum inhibits biofilm formation in *Staphylococcus aureus*. J. Microb. Pathog., 49 (6):388–391.
- ◆ Al Ahmary, M S.; Al Mastour, A S. and Ghnnam, W M. (2012). The Microbiology of tonsils in Khamis civil hospital, Saudi Arabia. ISRN Otolaryngology, (813581): 3.
- ◆ Bur, S.; Preissner, K T.; Herrmann, M. and Bischoff, M. (2013). The *Staphylococcus aureus* extracellular adherence protein promotes bacterial internalization by keratinocytes independent of fibronectin-binding proteins, J. Investig. Dermat., 133: 2004–2012 .
- ◆ Dakhil, B R. and Hamim, S S. (2016). Antibiotic susceptibility of *Streptococcus pyogenes* and *Staphylococcus aureus* isolated from pharyngitis and tonsillitis patients in Nasiriyah City. Iraq. World. J. Pharm. Sci., 4 (4): 14-19.
- ◆ Duran, N.; Dogramaci, Y.; Ozer, B.; Demir, C. and Kalaci, A. (2010). Detection of adhesin genes and slime production among Staphylococci in orthopaedic surgical wounds. J. Afric. Microb. Res., 4 (9): 708-715.
- ◆ Gowrishankar, S.; Thenmozhi, R.; Balaji, K. and Pandian, S K. (2013). Emergence of methicillin-resistant, vancomycin-intermediate

Staphylococcus aureus among patients associated with group A Streptococcal pharyngitis infection in southern India. J. Infec. Gene. Evol., 14 (1): 383–389.

◆ Gowrishankar, S.; Kamaladevi, A.; Balamurugan, K. and Pandian, S K. (2016). *In vitro* and *In vivo* biofilm characterization of methicillin resistant *Staphylococcus aureus* from patients associated with pharyngitis infection. J. Bio. Med. Research. International., Pp:1-14.

◆ Harley, J P. and Prescott, L M. (2002). Laboratory Exercises in Microbiology. (5th ed). The McGraw-Hill Companies, Inc., New York.

◆ Higgings, J.; loughman, A.; Van kessel, K P.; Van strijp, J A. and Foster, T J. (2006). Clumping Factor A of *Staphylococcus aureus* inhibits phagocytosis by human polymorphonuclear leucocytes. J. FEMS., 258 (2):290-296.

◆ Holban, A.; Cotar, A C.; Chifiriuc, M C.; Bleotu, C.; Banu, O. and Lazar, V. (2013). Variation of virulence profiles in some *Staphylococcus aureus* and *Pseudomonas aeruginosa* stains isolated from different clinical patients. J. Afric. Microb. Res., 7: 3453-3460.

◆ Jeong, J H.; Lee, DW.; Ryu, R A.; Lee, Y S.; Leeshh, L. and Kang, J O. (2007). Bacteriological comparison of tonsil core in recurrent tonsillitis and tonsillar hypertrophy. Laryngoscope, 117(12): 2146-2151.

◆ Kiavari, A.; Hasani, A.; Mobaiyen, H.; Aghazadeh, M.; Akbar Hasani, A.; Varshochi, M.; Rezaee, M A. and Dehghani, L. (2012). Detection of genes that encoding the collagen (Cna) and fibronectin (FnB) binding protein of *Staphylococcus aureus* isolates by multiplex- PCR. J. Med. Sci., 34 (4): 98-106.

◆ Lowy, F D. (2011). How *Staphylococcus aureus* adapts to its host. N. Engl. J. Med., 364:1987-1990.

◆ Mirzaee, M.; Peerayeha, S. N. and Behmanesh, M. (2015). Prevalence of fibronectin binding protein (*Fnb A* and *Fnb B*) genes among clinical isolates of methicillin resistant *Staphylococcus aureus*. J. Mol. Gene. Microb. Viro., 30 (4): 221–224.

◆ Nashev, D.; Toshkova, K.; Salasia, I O.; Hassan, A.; L€ammler, C. and Zsch€ock, M. (2004). Distribution of virulence genes of *Staphylococcus aureus* isolated from stable nasal carriers. J. FEMS. Microb. Lett., 233: 45–52.

- ◆ O'Neill, E.; Pozzi, C.; Houston, P.; Smyth, D.; Humphreys, H.; Robinson, D A. and O'Garra, J. (2007). Association between methicillin susceptibility and biofilm regulation in *Staphylococcus aureus* isolates from device-related infections. *J. Clin. Microbiol.*, 45: 1379-88.
- ◆ Peacock, S J.; Day, N P J.; Thomas, M G.; Berendt, A R. and Foster, T J. (2000). Clinical isolates of *Staphylococcus aureus* exhibit diversity in *fnb* genes and adhesion to human fibronectin. *J. Infect.*, 41 (1):23-31.
- ◆ Sabat, A.; Melles, D C.; Martirosian, G.; Grundmann, H.; Van belkum, A. and Hryniewicz, W. (2006). Distribution of the serine-aspartate repeat protein-encoding *sdr* genes among nasal-carriage and invasive *Staphylococcus aureus* strains. *J. Clin Microbiol.*, 44: 1135-1138.
- ◆ Sadoh, W. A.; Sadoh, A E.; Oladipo, A O. and Okunola, O O. (2008). Bacterial isolates of Tonsillitis and Pharyngitis in a Paediatric casualty setting. *J. Med. Biomed. Res.*, 7 (1): 38-44.
- ◆ Sivaraman, K.; Venkataraman, N. and Cole, A M. (2009). *Staphylococcus aureus* nasal carriage and its contributing factors. *J. Future Microbiol.*, 4: 999-1008.
- ◆ Tangchaisuriya, U.; Yotpanya, W.; Kittit, T. and Sitthisak, S. (2014). Distribution among Thai children of Methicillin-Resistant *Staphylococcus aureus* lacking *cna*, *fnbA* AND *icaAD*. *J. Southeast Asian Trop Med Public Health.* 45(1): 149-156.
- ◆ Thomas, R A. (2012). Comparative study of the core culture of tonsils in patients having recurrent/chronic tonsillitis vs. Tonsils with obstructive symptoms in tertiary care institute. Master Thesis. Pondicherry University.
- ◆ Wisniewska, K.; Garbacz, K. and Piechowicz, L. (2008). Genotypic screening of atypical *Staphylococcus aureus* strains isolated from clinical samples for presence of selected adhesin genes. *J. Med. Ma. Infect.*, 38: 549-53.
- ◆ Zautner, A E.; Krause, M.; Stropahl, G.; Holtfreter, S.; Frickmann, H.; Maletzki, C.; Kreikemeyer, B.; Hw, P. and Podbielski, A. (2010). Intracellular persisting *Staphylococcus aureus* is the major pathogen in recurrent tonsillitis. *J. PLoS. One.*, Pp:5.