

**Fuzzy neutrosophic Soft Ideal Theory Fuzzy neutrosophic
Soft Local Function and Generated Fuzzy neutrosophic
Soft Topological Spaces**

<https://doi.org/10.32792/utq/utj/vol14/3/2>

Ather Farhan lafta, Dr. Mohammed J. Al-Mesafer, Dr.

Ayed E. Hashoosh

College of Education for Pure Sciences,

University of Thi-Qar.

Atherfarhan2018@gmail.com Ayed197991@yahoo.com

Abstract: The purpose of this paper is to introduce the notion of fuzzy neutrosophic soft ideal in fuzzy neutrosophic soft set theory. The concept of fuzzy neutrosophic soft local function is also introduced. These concepts are discussed with a view to find new fuzzy neutrosophic soft topologies from the original one. The basic structure, especially a basis for such generated fuzzy neutrosophic soft topologies also studied here. Finally, the notion of compatibility of fuzzy neutrosophic soft ideals with fuzzy neutrosophic soft topologies is introduced and some equivalent conditions concerning this topic are established here.

Keywords: Soft set, Neutrosophic Soft, fuzzy neutrosophic Soft topological space , fuzzy neutrosophic Soft ideal, fuzzy neutrosophic Soft local function, *- fuzzy neutrosophic soft topology, Compatible fuzzy neutrosophic soft ideal.

1. Introduction:

The fuzzy set was presented by Zadeh [16] in 1965 where every element had a degree of membership. The intuitionistic fuzzy set on a universe X was presented by K. Atanaasov [3] in 1983 as a generalization of fuzzy set. The idea of Neutrosophic set was presented by F. Smarandache [14] which is a mathematical instrument for dealing with issues including uncertain, indeterminacy and conflicting information. In 1999, Molodtsov [10], initiated the novel concept of soft set theory, which was a totally new methodology for demonstrating

vulnerability and had a rich potential for application in a few ways is free from the troubles influencing existing techniques. P. K. Maji [9] had joined the Neutrosophic set with delicate sets and presented another scientific model 'Neutrosophic soft set'. A new notion of Fuzzy Neutrosophic soft set and its basic operations and results in Fuzzy Neutrosophic soft spaces are obtained.[1]. In 1966 Kuratowski [8] and Vaidyanathaswamy [15], introduced the concept of ideal in topological space and defined local function in ideal topological space. Further Hamlett and Jankovic in [6] studied the properties of ideal topological spaces.

Salama et al [11] defined intuitionistic fuzzy ideal for a set and generalized the concept of fuzzy ideal concepts, first initiated by Sarker [13]. Also, in 2014 A.A. Salama et al.[12] introduced and studied the concept of neutrosophic crisp local function, and construct a new type of neutrosophic crisp topological space via neutrosophic crisp ideals.

The purpose behind this paper is to present the idea of fuzzy neutrosophic soft ideal in fuzzy neutrosophic soft set theory which is the generalization of the concept of fuzzy intuitionistic ideal topological concepts and soft ideal topological concepts, first initiated by Salama [11], Kandil et al.[7] sequentially. The concept of fuzzy neutrosophic soft local function is also introduced. These ideas are talked about so as to discover new fuzzy neutrosophic soft topologies from the original one. The basic structure, especially a basis for such generated fuzzy neutrosophic soft topologies also studied here. At long last, the thought of compatibility of fuzzy neutrosophic soft ideals with fuzzy neutrosophic soft topologies is presented and some equivalent conditions concerning this theme are built up here.

2. Preliminaries

For the definitions and results on fuzzy neutrosophic soft set theory, we refer to [1,4,5]. However, we recall some definitions and results on fuzzy neutrosophic soft set theory and fuzzy neutrosophic soft topology.

Definition 2.1 [2]: A fuzzy neutrosophic set A on the universe X is defined as $A = \{ \langle x, T_A(x), I_A(x), F_A(x) \rangle; x \in X \}$ where the function $T, I, F: X \rightarrow [0,1]$. denote the degree of membership function (namely $T_A(x)$), the degree of indeterminacy function (namely $I_A(x)$) and the degree of non-membership (namely $F_A(x)$) respectively of each element $x \in X$ to the set A and

$$0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3$$

Definition 2.2 [10]: Let X be a universe set and E be a set of parameters. A pair (F, E) is called a soft set over X if and only if F is a function from E into the set of all subsets of X , i.e. $F: E \rightarrow P(X)$, where $P(X)$ is the power set of X . The set of all soft sets over X is denoted by $SS(X, E)$.

Definition 2.3 [1]: Let U be the initial universe set and E be a set of parameters. Consider a non-empty set A , $A \subseteq E$. Let $FNS(U)$ denote the set of all fuzzy neutrosophic sets of U . The collection $(N\tilde{F}, A)$ is termed to be the fuzzy neutrosophic soft set (FNS set or FNSS for short) over U , where $N\tilde{F}$ is a mapping given by $N\tilde{F}: A \rightarrow FNS(U)$. The family of all these fuzzy neutrosophic soft set over U with parameters from E denoted by $FNSS(U)_E$.

Definition 2.4 [1]: The complement of a fuzzy neutrosophic soft set $(N\tilde{F}, E)$ denoted by $(N\tilde{F}, A)^c$ and is defined as $(N\tilde{F}, A)^c = (N\tilde{F}^c, A)$ where $N\tilde{F}^c: A \rightarrow FNS(U)$ is a mapping given by $N\tilde{F}^c(e) = \{ \langle x, T_{N\tilde{F}^c(e)}(x) = F_{N\tilde{F}(e)}(x), I_{N\tilde{F}^c(e)}(x) = 1 - I_{N\tilde{F}(e)}(x), F_{N\tilde{F}^c(e)}(x) = T_{\tilde{F}(e)}(x) \rangle \}$.

Definition 2.5 [1]: A fuzzy neutrosophic soft set $(N\tilde{F}, E)$ over the universe U is said to be

(1) empty fuzzy neutrosophic soft set with respect to the parameter E if $T_{N\tilde{F}(e)} = 0, I_{N\tilde{F}(e)} = 0, F_{N\tilde{F}(e)} = 1, \forall x \in U, \forall e \in E$.

it is denoted by $\tilde{0}_E^U$.

(2) universe fuzzy neutrosophic soft set with respect to the parameter E if $T_{N\tilde{F}(e)} = 1, I_{N\tilde{F}(e)} = 1, F_{N\tilde{F}(e)} = 0, \forall x \in U, \forall e \in E$.

it is denoted by $\tilde{1}_E^U$.

Note: $(\tilde{0}_E^U)^c = \tilde{1}_E^U$ and $(\tilde{1}_E^U)^c = \tilde{0}_E^U$

Definition 2.6 [4]:

1. A fuzzy neutrosophic soft set $(N\tilde{F}, E)$ is said to be a fuzzy neutrosophic soft point, denoted by $e_{N\tilde{F}}$, if for the element $e \in E, N\tilde{F}(e) \neq \tilde{0}_N$ and $N\tilde{F}(e') = \tilde{0}_N, \forall e' \in E - \{e\}$. A class of all fuzzy neutrosophic soft points in U is denoted as $FNSP(U)_E$

2. The complement of a fuzzy neutrosophic soft point $e_{N\tilde{F}}$ is a fuzzy neutrosophic soft point $e_{N\tilde{F}}^c$ such that $N\tilde{F}^c(e) = (N\tilde{F}(e))^c$

3. A fuzzy neutrosophic soft point $e_{N\tilde{F}}$ is said to be in a fuzzy neutrosophic soft set $(N\tilde{G}, E)$ (or $e_{N\tilde{F}}$ is a fuzzy neutrosophic soft point of $(N\tilde{G}, E)$), denoted by $e_{N\tilde{F}} \in (N\tilde{G}, E)$ if for the element $e \in E, N\tilde{F}(e) \subseteq N\tilde{G}(e)$.

Example 2.7:

Let $U = \{x_1, x_2, x_3\}$ and $E = \{e_1, e_2\}$. Then,

$$e_{1N\tilde{F}} = \{ \langle x_1, (0.6, 0.4, 0.8) \rangle, \langle x_2, (0.8, 0.3, 0.5) \rangle, \langle x_3, (0.3, 0.7, 0.6) \rangle \}$$

is a fuzzy neutrosophic soft point whose complement is

$$e_{1N\tilde{F}}^c = \{ \langle x_1, (0.8, 0.6, 0.6) \rangle, \langle x_2, (0.5, 0.7, 0.8) \rangle, \langle x_3, (0.6, 0.3, 0.3) \rangle \}.$$

For another FNSS $(N\tilde{G}, E)$ defined on same (U, E) ,

let, $N\tilde{G}(e_1) = \{ \langle x_1, (0.7, 0.4, 0.7) \rangle, \langle x_2, (0.8, 0.5, 0.4) \rangle, \langle x_3, (0.5, 0.8, 0.5) \rangle \}$. Then, $N\tilde{F}(e_1) \subseteq N\tilde{G}(e_1)$ i.e., $e_{1N\tilde{F}} \in (N\tilde{G}, E)$.

Theorem 2.8: if $(N\tilde{F}, E)$ be a fuzzy neutrosophic soft set of U . implies $(N\tilde{F}, E) = \tilde{U}\{e_{N\tilde{F}} : e_{N\tilde{F}} \in (N\tilde{F}, E)\}$

Proof. Let $E = \{e_i : i \in I\}$, $(N\tilde{F}, E)$ be a fuzzy neutrosophic soft set and $\{e_{N\tilde{G}_k}\}_{k \in \Omega}$ be family of all fuzzy neutrosophic soft points of $(N\tilde{F}, E)$. For all $e_i \in E$, because of

$$N\tilde{F}(e_i) = \bigcup_{k \in \Omega} \tilde{G}_k(e_i)$$

$$\text{we have } (N\tilde{F}, E) = \{N\tilde{F}(e_i) : e_i \in E\} = \tilde{U}_{k \in \Omega} \tilde{G}_k.$$

Theorem 2.9: Let $(N\tilde{F}, E)$ and $(N\tilde{G}, E)$ be two fuzzy neutrosophic soft sets in $FNSS(U)_E$. Then $(N\tilde{F}, E) \cong (N\tilde{G}, E)$ if and only if $e_{N\tilde{H}} \in (N\tilde{F}, E)$ implies $e_{N\tilde{H}} \in (N\tilde{G}, E)$.

Proof. (\Rightarrow) : Let $(N\tilde{F}, E) \cong (N\tilde{G}, E)$. Therefore, for all $e \in E$, $N\tilde{F}(e) \cong N\tilde{G}(e)$. If $e_{N\tilde{H}} \in (N\tilde{F}, E)$, then $N\tilde{H}(e) \cong N\tilde{F}(e)$. Because of $N\tilde{H}(e) \cong N\tilde{F}(e) \cong N\tilde{G}(e)$, we have $N\tilde{H}(e) \cong N\tilde{G}(e)$ and so, $e_{N\tilde{H}} \in (N\tilde{G}, E)$.

(\Leftarrow) : If, for every $e_{N\tilde{H}} \in (N\tilde{F}, E)$ implies $e_{N\tilde{H}} \in (N\tilde{G}, E)$ implies, then in accordance with Theorem 2.10,

$$\tilde{U}_{e_{N\tilde{H}} \in (N\tilde{F}, E)} e_{N\tilde{H}} = (N\tilde{F}, E) \text{ and this implies } \tilde{U}_{e_{N\tilde{H}} \in (N\tilde{G}, E)} e_{N\tilde{H}} = (N\tilde{G}, E). \text{ So, } (N\tilde{F}, E) \cong (N\tilde{G}, E).$$

Definition 2.10 [5]: Let $e_{N\tilde{F}}$ and $e'_{N\tilde{G}}$ be two fuzzy neutrosophic soft points. For the fuzzy neutrosophic soft points $e_{N\tilde{F}}$ and $e'_{N\tilde{G}}$ over a common universe U , we say that the fuzzy neutrosophic soft points are distinct points if $e_{N\tilde{F}} \tilde{\cap} e'_{N\tilde{G}} = \tilde{0}_E^U$. It is clear that $e_{N\tilde{F}}$ and $e'_{N\tilde{G}}$ are distinct fuzzy neutrosophic soft points if and only if $e \neq e'$ or $\neq y$.

Definition 2.11 [1]: Let $(N\tilde{F}, E)$ be FNS set on (U, E) and τ be a collection of fuzzy neutrosophic soft subsets of $(N\tilde{F}, E)$. $(N\tilde{F}, E)$ is called fuzzy neutrosophic soft topology (FNST) if the following conditions hold.

(i) $\tilde{0}_E^U, \tilde{1}_E^U \in \tau$

(ii) $(N\tilde{F}, E), (N\tilde{G}, E) \in \tau$ implies $(N\tilde{F}, E) \tilde{\cap} (N\tilde{G}, E) \in \tau$

(iii) $\{(N\tilde{F}_\alpha, E) : \alpha \in \Gamma\} \in \tau$ implies $\tilde{\cup}\{(N\tilde{F}_\alpha, E) : \alpha \in \Gamma\} \in \tau$

The triplet (U, τ, E) is called an Fuzzy neutrosophic soft topological space (FNSTS) over U . Every member of τ is called an fuzzy neutrosophic soft open set in U . $(N\tilde{F}, E)$ is called an fuzzy neutrosophic soft closed set in U if $(N\tilde{F}, E) \in \tau^c$, where $\tau^c = \{(N\tilde{F}, E)^c : (N\tilde{F}, E) \in \tau\}$.

Definition 2.12: Let \hat{J} be a non-null collection of fuzzy neutrosophic soft sets over a universe U with a fixed set of parameters E is called a fuzzy neutrosophic soft ideal (FNSL for short) on U if

(1) $(N\tilde{F}, E) \in \hat{J}$ and $(N\tilde{G}, E) \in \hat{J} \Rightarrow (N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \in \hat{J}$ [Finite additivity],

(2) $(N\tilde{F}, E) \in \hat{J}$ and $(N\tilde{G}, E) \tilde{\subseteq} (N\tilde{F}, E) \Rightarrow (N\tilde{G}, E) \in \hat{J}$ [heredity].

If (U, τ, E) be a Fuzzy neutrosophic soft topological space, then (U, τ, E, \hat{J}) is called a Fuzzy neutrosophic soft ideal topological space (FNSITS, for short).

3. Fuzzy Neutrosophic Soft Local Function and Generated Fuzzy Neutrosophic Soft Topology

Definition 3.1: Let (U, τ, E) be a fuzzy neutrosophic soft topological space and \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E . $(N\tilde{F}, E)^*(\hat{J}, \tau)$ or $(N\tilde{F}, E)^* = \tilde{\cup}\{e_{N\tilde{F}} \in FN\tilde{S}P(U)_E : (N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J}, \forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau\}$

is called the fuzzy neutrosophic soft local function of $(N\tilde{F}, E)$ with respect to \hat{J} and τ , where $(N\tilde{O}, E)_{e_{N\tilde{F}}}$ is a τ -open fuzzy neutrosophic soft set containing $e_{N\tilde{F}}$.

Example 3.2:

(1) If $\hat{J} = \{ \tilde{0}_E^U \}$, then $(N\tilde{F}, E)^*(\hat{J}, \tau) = FNScl(N\tilde{F}, E)$ for all $(N\tilde{F}, E) \in FNSS(U)_E$.

(2) If $\hat{J} = FNSS(U)_E$, then $(N\tilde{F}, E)^*(\hat{J}, \tau) = \tilde{0}_E^U$ for all $(N\tilde{F}, E) \in FNSS(U)_E$.

Theorem 3.3: Let \hat{J} and \hat{J}_1 be any two fuzzy neutrosophic soft ideals with the same set of parameters E on a fuzzy neutrosophic soft topological space (U, τ, E) . Let $(N\tilde{F}, E), (N\tilde{G}, E) \in FNSS(U)_E$. Then

(1) $(\tilde{0}_E^U)^* = \tilde{0}_E^U$,

(2) $(N\tilde{F}, E) \subseteq (N\tilde{G}, E) \implies (N\tilde{F}, E)^* \subseteq (N\tilde{G}, E)^*$,

(3) $\hat{J} \subseteq \hat{J}_1 \implies (N\tilde{F}, E)^*(\hat{J}_1, \tau) \subseteq (N\tilde{F}, E)^*(\hat{J}, \tau)$,

(4) $(N\tilde{F}, E)^* = FNScl(N\tilde{F}, E)^* \subseteq FNScl(N\tilde{F}, E)$, where $FNScl$ is the fuzzy neutrosophic soft closure w.r.t. τ ,

(5) $(N\tilde{F}, E)^*$ is τ -closed fuzzy neutrosophic soft set.

(6) $((N\tilde{F}, E)^*)^* \subseteq (N\tilde{F}, E)^*$,

(7) $((N\tilde{F}, E) \cup (N\tilde{G}, E))^* = (N\tilde{F}, E)^* \cup (N\tilde{G}, E)^*$,

(8) $((N\tilde{F}, E) \cap (N\tilde{G}, E))^* \subseteq (N\tilde{F}, E)^* \cap (N\tilde{G}, E)^*$,

(9) $(N\tilde{H}, E) \in \hat{J} \implies ((N\tilde{F}, E) \cup (N\tilde{H}, E))^* = (N\tilde{F}, E)^*$

Proof.

(1) $(\tilde{0}_E^U)^* = \tilde{U}\{ e_{N\tilde{F}} \in \tilde{1}_E^U : (N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} \tilde{0}_E^U = \tilde{0}_E^U \in \hat{J} \forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau \} = \tilde{0}_E^U$.

(2) Let $e_{N\tilde{F}} \in (N\tilde{F}, E)^*$. Then $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J} \forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau$. Since $(N\tilde{F}, E) \subseteq (N\tilde{G}, E) \implies$

$(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \cong (N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{G}, E)$ and
 $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J}$. Then $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{G}, E) \notin \hat{J}$
 $\forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau$ from Definition 3.1. Hence $e_{N\tilde{F}} \in (N\tilde{G}, E)^*$. Thus
 $(N\tilde{F}, E)^* \cong (N\tilde{G}, E)^*$.

(3) Let $e_{N\tilde{F}} \in (N\tilde{F}, E)^*(\hat{J}_1, \tau)$. Then $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J}_1$
 $\forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau$. Since $\hat{J} \cong \hat{J}_1$. Then $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J}$
 $\forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau$. Hence $e_{N\tilde{F}} \in (N\tilde{F}, E)^*(\hat{J}, \tau)$. Thus
 $(N\tilde{F}, E)^*(\hat{J}_1, \tau) \cong (N\tilde{F}, E)^*(\hat{J}, \tau)$.

(4) since $\{\tilde{0}_E^U\} \cong \hat{J}$ for any FNS ideal on U, therefore by (3) and
 Example 3.2, $(N\tilde{F}, E)^*(\hat{J}) \cong (N\tilde{F}, E)^*(\tilde{0}_E^U) = FNScl(N\tilde{F}, E)$ for any
 FNSS $(N\tilde{F}, E)$ on U.

Suppose, $e_{N\tilde{F}} \in FNScl(N\tilde{F}, E)^*$. So for every $(N\tilde{O}, E)_{e_{N\tilde{F}}}$ open FNS set
 contains $e_{N\tilde{F}}$, $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E)^* \neq \tilde{0}_E^U$, there exists $e_{1N\tilde{G}} \in$
 $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E)^*$ such that for every $(N\tilde{V}, E)_{e_{1N\tilde{G}}}$ open FNS set
 contains $e_{1N\tilde{G}}$, $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J}$. Since
 $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{V}, E)_{e_{1N\tilde{G}}}$ open FNS set contains $e_{1N\tilde{G}}$ then
 $\left((N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{V}, E)_{e_{1N\tilde{G}}} \right) \tilde{\cap} (N\tilde{F}, E) \notin \hat{J}$ which leads to
 $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J} \forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau$, therefore $e_{N\tilde{F}} \in$
 $(N\tilde{F}, E)^*$, and so, $FNScl(N\tilde{F}, E)^* \cong (N\tilde{F}, E)^*$ and since
 $(N\tilde{F}, E)^* \cong FNScl(N\tilde{F}, E)^*$. Hence $(N\tilde{F}, E)^* = FNScl(N\tilde{F}, E)^*$.

(5) from (4) $(N\tilde{F}, E)^* = FNScl(N\tilde{F}, E)^*$ then $(N\tilde{F}, E)^*$ is τ -closed fuzzy
 neutrosophic soft set.

(6) By (4), we have $((N\tilde{F}, E)^*)^* =$
 $FNScl((N\tilde{F}, E)^*)^* \cong FNScl(N\tilde{F}, E)^* = (N\tilde{F}, E)^*$.

(7) Let $e_{N\tilde{F}} \in \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right)^*$. Then

$(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right) \notin \hat{J} \forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau$, i.e

$(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \tilde{\cup} (N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{G}, E) \notin \hat{J} \forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau$. then,

we have two cases $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J}$ and $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{G}, E) \in \hat{J}$

or the converse, if $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \in \hat{J}$ and $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{G}, E) \in \hat{J}$

,this means that $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \tilde{\cup} (N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{G}, E) \in \hat{J}$, which

contradicts the hypothesis. Hence

$$\left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right)^* \cong (N\tilde{F}, E)^* \tilde{\cup} (N\tilde{G}, E)^*.$$

For the reverse inclusion, since $(N\tilde{F}, E), (N\tilde{G}, E) \cong \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right)$. Then

$(N\tilde{F}, E)^* \cong \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right)^*$ and $(N\tilde{G}, E)^* \cong \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right)^*$ from

(2). Hence $(N\tilde{F}, E)^* \tilde{\cup} (N\tilde{G}, E)^* \cong \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right)^*$ and it implies that

$$\left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right)^* = (N\tilde{F}, E)^* \tilde{\cup} (N\tilde{G}, E)^*.$$

(8) Since $\left((N\tilde{F}, E) \tilde{\cap} (N\tilde{G}, E) \right) \cong (N\tilde{F}, E), (N\tilde{G}, E)$. Then

$\left((N\tilde{F}, E) \tilde{\cap} (N\tilde{G}, E) \right)^* \cong (N\tilde{F}, E)^*$ and

$\left((N\tilde{F}, E) \tilde{\cap} (N\tilde{G}, E) \right)^* \cong (N\tilde{G}, E)^*$ from (2). Hence

$$\left((N\tilde{F}, E) \tilde{\cap} (N\tilde{G}, E) \right)^* \cong (N\tilde{F}, E)^* \tilde{\cap} (N\tilde{G}, E)^*.$$

(9) Let $e_{N\tilde{F}} \in \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{H}, E) \right)^*$. Then

$(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{H}, E) \right) \notin \hat{J}, \forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau$, i.e

$(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \tilde{\cup} (N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{H}, E) \notin \hat{J}, \forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in$

τ . then, we have two cases $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \in \hat{J}$ and

$(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{H}, E) \notin \hat{J}$ or the converse, if $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{H}, E) \notin$

\hat{J}

gives $(N\tilde{H}, E) \notin \hat{J}$ which is a contradiction. Thus

$(N\tilde{O}, E)_{e_{N\tilde{F}}} \cap (N\tilde{F}, E) \notin \hat{J} \quad \forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau$. So $e_{N\tilde{F}} \in (N\tilde{F}, E)^*$

hence $\left((N\tilde{F}, E) \cup (N\tilde{H}, E) \right) \cong (N\tilde{F}, E)^*$.

For the reverse inclusion, since $(N\tilde{F}, E) \cong \left((N\tilde{F}, E) \cup (N\tilde{G}, E) \right)$. Then

$(N\tilde{F}, E)^* \cong \left((N\tilde{F}, E) \cup (N\tilde{G}, E) \right)^*$ from (2). Hence $\left((N\tilde{F}, E) \cup (N\tilde{H}, E) \right) = (N\tilde{F}, E)^*$.

Definition 3.4: Let (U, τ, E) be a fuzzy neutrosophic soft topological space and \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E . Then the operator $FNScl^*: FNSS(U)_E \rightarrow FNSS(U)_E$ defined by:
 $FNScl^*(N\tilde{F}, E) = (N\tilde{F}, E) \cup (N\tilde{F}, E)^*$ is a fuzzy neutrosophic soft closure operator.

Theorem 3.5:

(1) $FNScl^*(\tilde{O}_E^U) = (\tilde{O}_E^U)$.

(2) $(N\tilde{F}, E) \cong FNScl^*(N\tilde{F}, E), \forall (N\tilde{F}, E) \in FNSS(U)_E$.

(3) $FNScl^*\left((N\tilde{F}, E) \cup (N\tilde{G}, E) \right) = FNScl^*(N\tilde{F}, E) \cup FNScl^*(N\tilde{G}, E)$
 $, \forall (N\tilde{F}, E), (N\tilde{G}, E) \in FNSS(U)_E$.

(4) $FNScl^*\left(FNScl^*(N\tilde{F}, E) \right) \cong FNScl^*(N\tilde{F}, E), \forall (N\tilde{F}, E) \in FNSS(U)_E$.

Proof.

(1) since $FNScl^*(\tilde{O}_E^U) = (\tilde{O}_E^U) \cup (\tilde{O}_E^U)^* = \tilde{O}_E^U \cup \tilde{O}_E^U = \tilde{O}_E^U$ from Theorem 3.3.(1).

(2) since $(N\tilde{F}, E) \cong (N\tilde{F}, E) \cup (N\tilde{F}, E)^* = FNScl^*(N\tilde{F}, E)$

Thus $(N\tilde{F}, E) \cong FNScl^*(N\tilde{F}, E)$.

$$\begin{aligned}
 (3) \quad & FNScl^* \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right) = \\
 & \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right) \tilde{\cup} \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right)^* = \\
 & \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{G}, E) \right) \tilde{\cup} \left((N\tilde{F}, E)^* \tilde{\cup} (N\tilde{G}, E)^* \right) = \\
 & \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* \right) \tilde{\cup} \left((N\tilde{G}, E) \tilde{\cup} (N\tilde{G}, E)^* \right) = \\
 & FNScl^*(N\tilde{F}, E) \tilde{\cup} FNScl^*(N\tilde{G}, E) \text{ from Theorem 3.3. (7).}
 \end{aligned}$$

$$\begin{aligned}
 (4) \quad & FNScl^* \left(FNScl^*(N\tilde{F}, E) \right) = FNScl^* \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* \right) \\
 & = \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* \right) \tilde{\cup} \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* \right)^* = \\
 & \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* \right) \tilde{\cup} \left((N\tilde{F}, E)^* \tilde{\cup} ((N\tilde{F}, E)^*)^* \right) \cong \\
 & \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* \right) \tilde{\cup} \left((N\tilde{F}, E)^* \tilde{\cup} (N\tilde{F}, E)^* \right) = \\
 & \left((N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* \right) \\
 & = FNScl^*(N\tilde{F}, E) \text{ from Theorem 3.3. (6),(7) ,thus} \\
 & FNScl^* \left(FNScl^*(N\tilde{F}, E) \right) \cong FNScl^*(N\tilde{F}, E).
 \end{aligned}$$

Definition 3.6 : Let (U, τ, E) be a fuzzy neutrosophic soft topological space, \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E and $FNScl^*: FNSS(U)_E \rightarrow FNSS(U)_E$ be the fuzzy neutrosophic soft closure operator. Then there exists a unique fuzzy neutrosophic soft topology over U with the same set of parameters E , finer than τ , called the $*$ -fuzzy neutrosophic soft topology, denoted by $\tau^*(\hat{J})$ or τ^* , given by

$$\tau^*(\hat{J}) = \{ (N\tilde{F}, E) \in FNSS(U)_E : FNScl^*(N\tilde{F}, E)^c = (N\tilde{F}, E)^c \}.$$

Example 3.7: Let $U = \{r_1, r_2\}$, $E = \{e\}$ and $\tau = \{\tilde{1}_E^U, \tilde{0}_E^U, (N\tilde{F}, E)\}$ where $(N\tilde{F}, E)$ is a fuzzy neutrosophic soft set over U defined by

$N\tilde{F}(e) = \{ \langle r_1, (0.8, 0.7, 0.5) \rangle, \langle r_2, (0.9, 0.6, 0.5) \rangle \}$. Then τ defines a Fuzzy neutrosophic soft topology on U.

Let $\hat{J} = \{ \tilde{0}_E^U, (N\tilde{G}, E) \}$ be a fuzzy neutrosophic soft ideal over U where $(N\tilde{G}, E)$ is a fuzzy neutrosophic soft set over U defined by

$N\tilde{G}(e) = \{ \langle r_1, (0.8, 0.7, 0.5) \rangle, \langle r_2, (0.9, 0.6, 0.5) \rangle \}$. Then, $\tau^* = \{ \tilde{1}_E^U, \tilde{0}_E^U, (N\tilde{F}_1, E), (N\tilde{F}_2, E) \}$, where $(N\tilde{F}_1, E), (N\tilde{F}_2, E)$ are fuzzy neutrosophic soft sets over U where

$$N\tilde{F}_1(e) = \{ \langle r_1, (0.8, 0.7, 0.5) \rangle, \langle r_2, (0.9, 0.6, 0.5) \rangle \}$$

$$N\tilde{F}_2(e) = \{ \langle r_1, (0.5, 0.3, 0.8) \rangle, \langle r_2, (0.5, 0.4, 0.9) \rangle \}.$$

Example 3.8:

(1) If $\hat{J} = \{ \tilde{0}_E^U \}$, then $FNScl^*(N\tilde{F}, E) = FNScl(N\tilde{F}, E)$ for all $(N\tilde{F}, E) \in FNSS(U)_E$ and $\tau^* = \tau$.

(2) If $\hat{J} = FNSS(U)_E$, then $FNScl^*(N\tilde{F}, E) = (N\tilde{F}, E)$ for all $(N\tilde{F}, E) \in FNSS(U)_E$ and $\tau^* = FNSS(U)_E$ (the fuzzy neutrosophic soft discrete topology).

Proof.

(1) Let $\hat{J} = \{ \tilde{0}_E^U \} \implies (N\tilde{F}, E)^*(\hat{J}, \tau) = FNScl(N\tilde{F}, E)$ from Example 3.2

$$\text{And since } FNScl^*(N\tilde{F}, E) = (N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* = (N\tilde{F}, E) \tilde{\cup} FNScl(N\tilde{F}, E)$$

Since $(N\tilde{F}, E) \tilde{\subseteq} FNScl(N\tilde{F}, E)$ then $(N\tilde{F}, E) \tilde{\cup} FNScl(N\tilde{F}, E) = FNScl(N\tilde{F}, E)$, Thus $FNScl^*(N\tilde{F}, E) = FNScl(N\tilde{F}, E)$.

Now, let $(N\tilde{F}, E) \in \tau^*$, $FNScl^*(N\tilde{F}, E) = FNScl(N\tilde{F}, E)$

$$\text{Since } (N\tilde{F}, E) \in \tau^* \implies FNScl^*(N\tilde{F}, E)^c = (N\tilde{F}, E)^c$$

So, $FNScl^*(N\tilde{F}, E)^c = FNScl(N\tilde{F}, E)^c = (N\tilde{F}, E)^c$

Thus $(N\tilde{F}, E)^c$ is closed FNS set $\Rightarrow (N\tilde{F}, E)$ is open FNS set i.e $(N\tilde{F}, E) \in \tau$

Hence $\tau^* \cong \tau$. But from Definition 3.6. $\tau \cong \tau^*$, hence $\tau^* = \tau$.

(2) Let $\hat{J} = FNSS(U)_E \Rightarrow (N\tilde{F}, E)^*(\hat{J}, \tau) = \tilde{0}_E^U$ from Example 3.3.
since $FNScl^*(N\tilde{F}, E) = (N\tilde{F}, E) \cup (N\tilde{F}, E)^* = (N\tilde{F}, E) \cup \tilde{0}_E^U = (N\tilde{F}, E)$

So, $FNScl^*(N\tilde{F}, E) = (N\tilde{F}, E)$.

Now, let $(N\tilde{F}, E) \in FNSS(U)_E$, $FNScl^*(N\tilde{F}, E) = (N\tilde{F}, E)$ So,
 $FNScl^*(N\tilde{F}, E)^c = (N\tilde{F}, E)^c$, thus $(N\tilde{F}, E) \in \tau^*$

Hence $FNSS(U)_E \cong \tau^*$, and from Definition 3.6. $\tau^* \cong FNSS(U)_E$

Thus $\tau^* = FNSS(U)_E$.

Theorem 3.9: Let (U, τ, E) be a fuzzy neutrosophic soft topological space and \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E. Then

$\sigma(\hat{J}, \tau) = \{(N\tilde{F}, E) - (N\tilde{G}, E) : (N\tilde{F}, E) \in \tau, (N\tilde{G}, E) \in \hat{J}\}$ forms a basis for the generated fuzzy neutrosophic soft topology $\tau^*(\hat{J})$.

Proof. Since $\tilde{1}_E^U \in \tau$, $\tilde{0}_E^U \in \hat{J}$. Then $\tilde{1}_E^U - \tilde{0}_E^U \in \sigma$. Hence $\tilde{1}_E^U \in \sigma$ and $\bigcup_{j \in J} ((N\tilde{F}_j, E) - (N\tilde{G}_j, E)) = \tilde{1}_E^U$.

Also, let $((N\tilde{F}_1, E) - (N\tilde{G}_1, E)), ((N\tilde{F}_2, E) - (N\tilde{G}_2, E)) \in \sigma$ such that

$e_{N\tilde{F}} \in ((N\tilde{F}_1, E) - (N\tilde{G}_1, E)) \tilde{\cap} ((N\tilde{F}_2, E) - (N\tilde{G}_2, E))$. Then

$e_{N\tilde{F}} \in ((N\tilde{F}_1, E) - (N\tilde{G}_1, E)) \tilde{\cap} ((N\tilde{F}_2, E) - (N\tilde{G}_2, E))$

$$\begin{aligned}
 &= \left((N\tilde{F}_1, E) \tilde{\cap} (N\tilde{G}_1, E)^c \right) \tilde{\cap} \left((N\tilde{F}_2, E) \tilde{\cap} (N\tilde{G}_2, E)^c \right) \\
 &= \left((N\tilde{F}_1, E) \tilde{\cap} (N\tilde{F}_2, E) \right) \tilde{\cap} \left((N\tilde{G}_1, E)^c \tilde{\cap} (N\tilde{G}_2, E)^c \right) \\
 &= \left((N\tilde{F}_1, E) \tilde{\cap} (N\tilde{F}_2, E) \right) \tilde{\cap} \left((N\tilde{G}_1, E) \tilde{\cup} (N\tilde{G}_2, E) \right)^c \\
 &= \left((N\tilde{F}_1, E) \tilde{\cap} (N\tilde{F}_2, E) \right) - \left((N\tilde{G}_1, E) \tilde{\cap} (N\tilde{G}_2, E) \right) \in \sigma(\hat{J}, \tau)
 \end{aligned}$$

Thus σ is a fuzzy neutrosophic soft basis of τ^* .

Corollary 3.10: Let (U, τ, E) be a fuzzy neutrosophic soft topological space and \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E . Then $\tilde{\subseteq} \sigma(\hat{J}, \tau) \tilde{\subseteq} \tau^*(\hat{J})$.

Proof. Let $(N\tilde{F}, E) \in \tau$, since $\tilde{0}_E^U \in \hat{J}$

Thus $(N\tilde{F}, E) = (N\tilde{F}, E) - \tilde{0}_E^U \in \sigma(\hat{J}, \tau)$ i.e $(N\tilde{F}, E) \in \sigma(\hat{J}, \tau)$

So, $\tau \tilde{\subseteq} \sigma(\hat{J}, \tau)$, and since $\sigma(\hat{J}, \tau)$ is a fuzzy neutrosophic soft basis of $\tau^*(\hat{J})$, Then $\sigma(\hat{J}, \tau) \tilde{\subseteq} \tau^*(\hat{J})$, hence $\tau \tilde{\subseteq} \sigma(\hat{J}, \tau) \tilde{\subseteq} \tau^*(\hat{J})$.

Theorem 3.11: Let \hat{J}_1, \hat{J}_2 are a fuzzy neutrosophic soft ideals over U , $\hat{J}_1 \tilde{\subseteq} \hat{J}_2$ then $\tau^*(\hat{J}_1) \tilde{\subseteq} \tau^*(\hat{J}_2)$.

Proof. Let $(N\tilde{F}, E) \in \tau^*(\hat{J}_1) \Rightarrow FNScl^*(N\tilde{F}, E)^c = (N\tilde{F}, E)^c$

$\Rightarrow FNScl^*(N\tilde{F}, E)^c = (N\tilde{F}, E)^c \tilde{\cup} ((N\tilde{F}, E)^c)^*(\hat{J}_1, \tau) = (N\tilde{F}, E)^c$ from definition 3.6.

$\Rightarrow [(N\tilde{F}, E)^c]^c = \left[(N\tilde{F}, E)^c \tilde{\cup} ((N\tilde{F}, E)^c)^*(\hat{J}_1, \tau) \right]^c$.

$\Rightarrow (N\tilde{F}, E) = (N\tilde{F}, E) \tilde{\cap} \left[((N\tilde{F}, E)^c)^*(\hat{J}_1, \tau) \right]^c$

Since $\hat{J}_1 \cong \hat{J}_2$ then $((N\tilde{F}, E)^c)^* (\hat{J}_2, \tau) \cong ((N\tilde{F}, E)^c)^* (\hat{J}_1, \tau)$ from Theorem 3.3. (3)

$$\Rightarrow \left[((N\tilde{F}, E)^c)^* \right]^c (\hat{J}_1, \tau) \cong \left[((N\tilde{F}, E)^c)^* \right]^c (\hat{J}_2, \tau)$$

$$\Rightarrow (N\tilde{F}, E) =$$

$$(N\tilde{F}, E) \tilde{\cap} \left[((N\tilde{F}, E)^c)^* \right]^c (\hat{J}_1, \tau) \cong (N\tilde{F}, E) \tilde{\cap} \left[((N\tilde{F}, E)^c)^* \right]^c (\hat{J}_2, \tau)$$

$$\text{But } (N\tilde{F}, E) \tilde{\cap} \left[((N\tilde{F}, E)^c)^* \right]^c (\hat{J}_2, \tau) \cong (N\tilde{F}, E), \text{ thus } (N\tilde{F}, E) =$$

$$(N\tilde{F}, E) \tilde{\cap} \left[((N\tilde{F}, E)^c)^* \right]^c (\hat{J}_2, \tau) \Rightarrow (N\tilde{F}, E)^c \tilde{\cup} ((N\tilde{F}, E)^c)^* (\hat{J}_2, \tau) = (N\tilde{F}, E)^c$$

Thus, $FNScl^*(N\tilde{F}, E)^c = (N\tilde{F}, E)^c$ i. e $(N\tilde{F}, E) \in \tau^*(\hat{J}_2)$, hence $\tau^*(\hat{J}_1) \cong \tau^*(\hat{J}_2)$.

Theorem 3.12: Let τ_1, τ_2 be two fuzzy neutrosophic soft topologies on U. Then for any fuzzy neutrosophic soft ideal \hat{J} on U, $\tau_1 \cong \tau_2$ implies $(N\tilde{F}, E)^*(\hat{J}, \tau_2) \cong (N\tilde{F}, E)^*(\hat{J}, \tau_1)$, $\forall (N\tilde{F}, E) \in \hat{J}$ and $\tau_1^* \cong \tau_2^*$

Proof. Let $e_{N\tilde{F}} \in (N\tilde{F}, E)^*(\hat{J}, \tau_2) \Rightarrow (N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J}$
 $\forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau_2$ Since $\tau_1 \cong \tau_2 \Rightarrow (N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \notin \hat{J}$
 $\forall (N\tilde{O}, E)_{e_{N\tilde{F}}} \in \tau_2$, Thus $e_{N\tilde{F}} \in (N\tilde{F}, E)^*(\hat{J}, \tau_1)$, hence $(N\tilde{F}, E)^*(\hat{J}, \tau_2) \cong (N\tilde{F}, E)^*(\hat{J}, \tau_1)$.

Next, we proved $\tau_1^* \cong \tau_2^*$, Let $(N\tilde{F}, E) \in \tau_1^* \Rightarrow FNScl_{\tau_1}^*(N\tilde{F}, E)^c = (N\tilde{F}, E)^c$

$\Rightarrow FNScl_{\tau_1}^*(N\tilde{F}, E)^c = (N\tilde{F}, E)^c \tilde{\cup} ((N\tilde{F}, E)^c)^*(\hat{J}, \tau_1) = (N\tilde{F}, E)^c$ from definition 3.6.

$$\Rightarrow ((N\tilde{F}, E)^c)^c = \left((N\tilde{F}, E)^c \tilde{\cup} ((N\tilde{F}, E)^c)^*(\hat{J}, \tau_1) \right)^c$$

$$\Rightarrow (N\tilde{F}, E) = (N\tilde{F}, E) \tilde{\cap} \left[\left((N\tilde{F}, E)^c \right)^* \right]^c (\hat{J}, \tau_1)$$

Since $\tau_1 \cong \tau_2$ then $\left((N\tilde{F}, E)^c \right)^* (\hat{J}, \tau_2) \cong \left((N\tilde{F}, E)^c \right)^* (\hat{J}, \tau_1)$ from above,

$$\Rightarrow \left[\left((N\tilde{F}, E)^c \right)^* \right]^c (\hat{J}, \tau_1) \cong \left[\left((N\tilde{F}, E)^c \right)^* \right]^c (\hat{J}, \tau_2),$$

$$\Rightarrow (N\tilde{F}, E) = (N\tilde{F}, E) \tilde{\cap} \left[\left((N\tilde{F}, E)^c \right)^* \right]^c (\hat{J}, \tau_1) \cong$$

$$(N\tilde{F}, E) \tilde{\cap} \left[\left((N\tilde{F}, E)^c \right)^* \right]^c (\hat{J}, \tau_2), \text{ But } (N\tilde{F}, E) \tilde{\cap} \left[\left((N\tilde{F}, E)^c \right)^* \right]^c (\hat{J}, \tau_2)$$

$$\cong (N\tilde{F}, E), \text{ so, } (N\tilde{F}, E) = (N\tilde{F}, E) \tilde{\cap} \left[\left((N\tilde{F}, E)^c \right)^* \right]^c (\hat{J}, \tau_2),$$

$$\Rightarrow (N\tilde{F}, E)^c \tilde{\cup} \left((N\tilde{F}, E)^c \right)^* (\hat{J}, \tau_2) = (N\tilde{F}, E)^c$$

Thus, $FNScl_{\tau_2}^*(N\tilde{F}, E)^c = (N\tilde{F}, E)^c$ i.e $(N\tilde{F}, E) \in \tau_2^*$.

4. Compatibility of Fuzzy Neutrosophic Soft Ideals with Fuzzy Neutrosophic Soft Topology

Definition 4.1: Let (U, τ, E) be a fuzzy neutrosophic soft topological space and \hat{J} be a fuzzy neutrosophic soft ideal on U . τ is said to be compatible with \hat{J} , denoted by $\tau \approx \hat{J}$, if for every fuzzy neutrosophic soft set $(N\tilde{F}, E)$ of U and for all fuzzy neutrosophic soft point $e_{N\tilde{F}} \in (N\tilde{F}, E)$, there exists $(N\tilde{O}, E)_{e_{N\tilde{F}}}$ open fuzzy neutrosophic soft contains $e_{N\tilde{F}}$ such that $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \in \hat{J}$, then $(N\tilde{F}, E) \in \hat{J}$.

Theorem 4.2: Let (U, τ, E) be a fuzzy neutrosophic soft topological space, \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E and $\tau \approx \hat{J}$. Then For any fuzzy neutrosophic soft set $(N\tilde{F}, E)$ of U , $(N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* = \tilde{O}_E^U$ implies $(N\tilde{F}, E)^* = \tilde{O}_E^U$ iff $(N\tilde{F}, E)^* = \left((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* \right)^*$.

Proof. (\Rightarrow) Suppose that $(N\tilde{F}, E)$ any fuzzy neutrosophic soft set of U ,

$$(N\tilde{F}, E)^* = \left((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* \right)^* \text{ and } (N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* = \tilde{0}_E^U.$$

Then $(N\tilde{F}, E)^* = \left((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* \right)^* = (\tilde{0}_E^U)^* = \tilde{0}_E^U$. Hence $(N\tilde{F}, E)^* = \tilde{0}_E^U$.

(\Leftarrow) Suppose that $(N\tilde{F}, E)$ any fuzzy neutrosophic soft set of U , $(N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* = \tilde{0}_E^U$ and $(N\tilde{F}, E)^* = \tilde{0}_E^U$. Since $(N\tilde{F}, E)^* = \tilde{0}_E^U$ and $(N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* = \tilde{0}_E^U \Rightarrow \left((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* \right)^* = (\tilde{0}_E^U)^* = \tilde{0}_E^U$, Hence $(N\tilde{F}, E)^* = \left((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* \right)^*$.

Corollary 4.3: Let (U, τ, E) be a fuzzy neutrosophic soft topological space, \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E , For any fuzzy neutrosophic soft set $(N\tilde{F}, E)$ of U and $\tau \approx \hat{J}$. Then $(N\tilde{F}, E)^* = \left((N\tilde{F}, E)^* \right)^*$.

Proof. Suppose that $(N\tilde{F}, E)$ any fuzzy neutrosophic soft set of U . Since $(N\tilde{F}, E)^* = \left((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* \right)^* \cong (N\tilde{F}, E)^* \tilde{\cap} \left((N\tilde{F}, E)^* \right)^* = \left((N\tilde{F}, E)^* \right)^*$ from Theorem 3.3. (8), (6). thus $(N\tilde{F}, E)^* = \left((N\tilde{F}, E)^* \right)^*$.

Theorem 4.4: Let (U, τ, E) be a fuzzy neutrosophic soft topological space, \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E . Then the following are equivalent:

- (a) $\tau \approx \hat{J}$.
- (b) For every $(N\tilde{F}, E) \in FNSS(U)_E$ such that $(N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* = \tilde{0}_E^U$. then $(N\tilde{F}, E) \in \hat{J}$.
- (c) For every $(N\tilde{F}, E) \in FNSS(U)_E$, $(N\tilde{F}, E) - (N\tilde{F}, E)^* \in \hat{J}$.

Proof. (a) \Rightarrow (b) Let $(N\tilde{F}, E) \in FNSS(U)_E$ such that $(N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* = \tilde{0}_E^U$, so, $(N\tilde{F}, E) \cong \left((N\tilde{F}, E)^* \right)^c$

Since $(N\tilde{F}, E)^*$ is τ -closed fuzzy neutrosophic soft set , Then for all $e_{N\tilde{F}} \in (N\tilde{F}, E)$ and $e_{N\tilde{F}} \notin (N\tilde{F}, E)^*$ such that $e_{N\tilde{F}} \in ((N\tilde{F}, E)^*)^c$, we have $((N\tilde{F}, E)^*)^c \tilde{\cap} (N\tilde{F}, E) \in \hat{J}$ for some $((N\tilde{F}, E)^*)^c \in \tau$. Thus $(N\tilde{F}, E) \in \hat{J}$ by (a).

$$\begin{aligned} (b) \Rightarrow (c) \text{ Let } (N\tilde{F}, E) \in FNSS(U)_E. \text{ Since} \\ (N\tilde{F}, E) - (N\tilde{F}, E)^* \tilde{\cap} ((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^*)^* = \\ ((N\tilde{F}, E) \tilde{\cap} ((N\tilde{F}, E)^*)^c) \tilde{\cap} ((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^*)^* \cong \\ ((N\tilde{F}, E) \tilde{\cap} ((N\tilde{F}, E)^*)^c) \tilde{\cap} (N\tilde{F}, E)^* \tilde{\cap} ((N\tilde{F}, E)^*)^* \\ \cong ((N\tilde{F}, E) \tilde{\cap} ((N\tilde{F}, E)^*)^c) \tilde{\cap} (N\tilde{F}, E)^* = \tilde{0}_E^U \end{aligned}$$

Then $(N\tilde{F}, E) - (N\tilde{F}, E)^* \in \hat{J}$ by (b).

$e_{N\tilde{F}} \in (N\tilde{F}, E)$, $((N\tilde{F}, E)^*)^c \tilde{\cap} (N\tilde{F}, E) \in \hat{J}$ such that $e_{N\tilde{F}} \in ((N\tilde{F}, E)^*)^c$

$\Rightarrow (N\tilde{F}, E) \cong ((N\tilde{F}, E)^*)^c$. Since $(N\tilde{F}, E)^*$ is τ -closed fuzzy neutrosophic soft set $\Rightarrow ((N\tilde{F}, E)^*)^c \in \tau$. Thus $(N\tilde{F}, E) \in \hat{J}$ by (b).

(c) \Rightarrow (a) suppose that $(N\tilde{F}, E) \in FNSS(U)_E$ and assume that for every $e_{N\tilde{F}} \in (N\tilde{F}, E)$ there exists $(N\tilde{O}, E)_{e_{N\tilde{F}}}$ open fuzzy neutrosophic soft contains $e_{N\tilde{F}}$ such that $(N\tilde{O}, E)_{e_{N\tilde{F}}} \tilde{\cap} (N\tilde{F}, E) \in \hat{J}$ and $(N\tilde{F}, E) - (N\tilde{F}, E)^* \in \hat{J}$. Then $e_{N\tilde{F}} \notin (N\tilde{F}, E)^*$ such that $(N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* = \tilde{0}_E^U$. Hence $(N\tilde{F}, E) \cong ((N\tilde{F}, E)^*)^c$. Since $(N\tilde{F}, E)^*$ is τ -closed fuzzy neutrosophic soft set $((N\tilde{F}, E)^*)^c \in \tau$ such that $e_{N\tilde{F}} \in ((N\tilde{F}, E)^*)^c$. Let $((N\tilde{F}, E)^*)^c = (N\tilde{O}, E)_{e_{N\tilde{F}}}$ such that $e_{N\tilde{F}} \in ((N\tilde{F}, E)^*)^c$. thus $((N\tilde{F}, E)^*)^c \tilde{\cap} (N\tilde{F}, E) \in \hat{J}$. Since

$(N\tilde{F}, E) \subseteq ((N\tilde{F}, E)^*)^c \implies (N\tilde{F}, E) = (N\tilde{F}, E) \tilde{\cap} ((N\tilde{F}, E)^*)^c = (N\tilde{F}, E) - (N\tilde{F}, E)^*$. by hypothesis, $(N\tilde{F}, E) - (N\tilde{F}, E)^* \in \hat{J}$. Thus $(N\tilde{F}, E) \in \hat{J}$. hence $\tau \approx \hat{J}$.

Theorem 4.5: Let (U, τ, E) be a fuzzy neutrosophic soft topological space, \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E . then For every $(N\tilde{F}, E) \in FNSS(U)_E$, $(N\tilde{F}, E) - (N\tilde{F}, E)^* \in \hat{J}$ if and only if $(N\tilde{F}, E)$ contains no non-null soft set $(N\tilde{G}, E)$ with $(N\tilde{G}, E) \subseteq (N\tilde{G}, E)^*$, then $(N\tilde{F}, E) \in \hat{J}$.

Proof. (\implies) Suppose that $(N\tilde{F}, E) \in FNSS(U)_E$ such that $(N\tilde{F}, E)$ contains no non-null fuzzy neutrosophic soft set (\tilde{G}, E) with $(N\tilde{G}, E) \subseteq (N\tilde{G}, E)^*$ and $(N\tilde{F}, E) - (N\tilde{F}, E)^* \in \hat{J}$. Since $((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^*) \subseteq (N\tilde{F}, E)^* = ((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^*)^*$ from Theorem(2.3.2). It follows that $(N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* \subseteq ((N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^*)^*$. By assumption, $(N\tilde{F}, E) \tilde{\cap} (N\tilde{F}, E)^* = \tilde{0}_E^U$, Thus $(N\tilde{F}, E) = (N\tilde{F}, E) - (N\tilde{F}, E)^* \in \hat{J}$.

(\impliedby) Suppose that $(N\tilde{F}, E) \in FNSS(U)_E$ such that $(N\tilde{F}, E)$ contains no non-null fuzzy neutrosophic soft set $(N\tilde{G}, E)$ with $(N\tilde{G}, E) \subseteq (N\tilde{G}, E)^*$ and $(N\tilde{F}, E) \in \hat{J}$. Since

$((N\tilde{F}, E) - (N\tilde{F}, E)^*) \tilde{\cap} ((N\tilde{F}, E) - (N\tilde{F}, E)^*)^* = \tilde{0}_E^U$ and $(N\tilde{F}, E) - (N\tilde{F}, E)^*$ contains no non-null fuzzy neutrosophic soft set $(N\tilde{G}, E)$ with $(N\tilde{G}, E) \subseteq (N\tilde{G}, E)^*$. Hence $(N\tilde{F}, E) - (N\tilde{F}, E)^* \in \hat{J}$.

Theorem 4.6: If (U, τ, E) is a fuzzy neutrosophic soft topological space, \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E and compatible with τ . Then a fuzzy neutrosophic soft set is τ^* -closed if and only if it is the union of a τ -closed fuzzy neutrosophic soft set and a fuzzy neutrosophic soft set in \hat{J} .

Proof. Let $(N\tilde{F}, E)$ be a τ^* -closed fuzzy neutrosophic soft set. Then $FNScl^*(N\tilde{F}, E) = (N\tilde{F}, E)$ and $(N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* = (N\tilde{F}, E)$. Hence $(N\tilde{F}, E)^* \cong (N\tilde{F}, E)$. Thus $(N\tilde{F}, E) = \left((N\tilde{F}, E) - (N\tilde{F}, E)^* \right) \tilde{\cup} (N\tilde{F}, E)^*$, $(N\tilde{F}, E) - (N\tilde{F}, E)^* \in \hat{J}$ from Theorem 4.4. and $(N\tilde{F}, E)^*$ is τ -closed fuzzy neutrosophic soft set from Theorem 3.3. (5).

Conversely, let $(N\tilde{F}, E) = (N\tilde{G}, E) \tilde{\cup} (NI, E)$, where $(N\tilde{G}, E)$ is τ -closed fuzzy neutrosophic soft set and $(NI, E) \in \hat{J}$. Then $(N\tilde{F}, E)^* = \left((N\tilde{G}, E) \tilde{\cup} (NI, E) \right)^* = (N\tilde{G}, E)^* \cong FNScl(N\tilde{G}, E) = (N\tilde{G}, E) \cong (N\tilde{F}, E)$ from Theorem 3.3. (2), (4). Hence $(N\tilde{F}, E) \tilde{\cup} (N\tilde{F}, E)^* = (N\tilde{F}, E)$. Thus $FNScl^*(N\tilde{F}, E) = (N\tilde{F}, E)$. It follows that $(N\tilde{F}, E)$ is a τ^* -closed fuzzy neutrosophic soft set.

Theorem 4.7: Let (U, τ, E) be a fuzzy neutrosophic soft topological space, \hat{J} be a fuzzy neutrosophic soft ideal over U with the same set of parameters E . If $\tau \approx \hat{J}$, then $\sigma(\hat{J}, \tau)$ is a fuzzy neutrosophic soft topology and hence $\sigma = \tau^*$.

Proof. Let $(N\tilde{G}, E) \in \tau^*$. Then $(N\tilde{G}, E)^c = (N\tilde{F}, E) \tilde{\cup} (NI, E)$, where $(N\tilde{F}, E)$ is τ -closed fuzzy neutrosophic soft set and $(NI, E) \in \hat{J}$. Hence $(N\tilde{G}, E) = \tilde{1}_E^U - \left((N\tilde{F}, E) \tilde{\cup} (NI, E) \right) = \left(\tilde{1}_E^U - (N\tilde{F}, E) \right) \tilde{\cap} \left(\tilde{1}_E^U - (NI, E) \right) = \left(\tilde{1}_E^U - (N\tilde{F}, E) \right) - (NI, E)$

, where $(N\tilde{F}, E) \in \tau$ and $(NI, E) \in \hat{J}$. Thus

$$(N\tilde{G}, E) = \left(\tilde{1}_E^U - (N\tilde{F}, E) \right) - (NI, E) \in \sigma(\hat{J}, \tau). \text{ So } \tau^* \cong \sigma(\hat{J}, \tau).$$

And since $\sigma(\hat{J}, \tau) \cong \tau^*$ from Corollary (2.2.11) Hence $\sigma = \tau^*$.

References

- [1] I. Arockiarani, I. R. Sumathi & J. Martina Jency -Fuzzy Neutrosophic soft topological spaces IJMA – 4 (10) 2013., 225-238

- [2] I. Arockiarani and J. Martina Jency: More on Fuzzy Neutrosophic Sets and Fuzzy Neutrosophic Topological Spaces, IJIRS, vol. 3, 2014, pp. 642-652.
- [3] K. Atanassov, Intuitionistic fuzzy sets, in V. Sgurev, ed., vii ITKRS Session, Sofia (June 1983 central Sci. and Techn. Library, Bulg. Academy of Sciences (1983)).
- [4] T. Bera and N. K. Mahapatra, Introduction to neutrosophic soft topological spaces, OPSEARCH, (March, 2017), DOI 10.1007/s12597-017-0308-7.
- [5] Gunduz C. and Taha Yasin ÖZTÜRK and Bayramov S, Separation axioms on neutrosophic soft topological spaces, Turk J Math, (2019) 43: 498-510, doi:10.3906/mat-1805-110.
- [6] T. R. Hamlett and D. Jankovic , Ideals in topological spaces and the set operator Ψ , Bull .M. U.M.I. , (7) , 4-B(1990) , 863-874.
- [7] A. Kandil, O. A. E. Tantaw, S. A. El-Sheikh and A. M. Abd El-latif, Soft ideal theory, Soft local function and generated soft topological spaces, Appl. Math. Inf. Sci. 8 (4) (2014) 1-9.
- [8] K. Kuratowski , Topology , Vol. I ,Academic press , Newyork , 1966 . [7] P. K.
- [9] Maji, Neutrosophic soft set, Annals of Fuzzy Mathematics and Informatics, Volume 5, No.1, (2013),157-168.
- [10] D. Molodtsov, Soft set Theory - First Results, Comput. Math. Appl. 37 (1999)19-31.
- [11] A. A. Salama and S.A. Alblowi, Intuitionistic Fuzzy Ideals Topological Spaces, Advances in Fuzzy Mathematics, Vol. (7), Number 1, pp. 51- 60, (2012).
- [12] A.A. Salama, F. Smarandache and S. A. Alblowi, New Neutrosophic Crisp Topological Concepts, Neutrosophic Sets and Systems, Vol. 4, 2014.

- [13] Debasis Sarker, Fuzzy ideal theory, Fuzzy local function and generated fuzzy topology, Fuzzy Sets and Systems 87, 117 – 123. (1997).
- [14] F. Smarandache, Neutrosophy, Neutrosophic Probability, Set and Logic, Amer. Res. Press, Rehoboth, USA., (1998), p. 105, <http://fs.gallup.unm.edu/eBook-neutrosophics4.pdf> (fourth version).
- [15] R. Vaidyanathaswamy , The localization theory in set topology . proc. Indian Acad. Sci. 20(1945) , 51-61
- [16] L. A. Zadeh, Fuzzy Sets, Inform and Control 8(1965) 338 – 353.